
Assessing how disruption of the methanogenic community and their syntrophic relationships in tidal freshwater marshes via saltwater intrusion may affect CH₄ emissions

A Data management plan created using the DMPTool

Creator(s): David Berrier

Affiliation: Virginia Commonwealth University

Last modified: July 02, 2015

Copyright information: The above plan creator(s) have agreed that others may use as much of the text of this plan as they would like in their own plans, and customize it as necessary. You do not need to credit the creators as the source of the language used, but using any of their plan's text does not imply that the creator(s) endorse, or have any relationship to, your project or proposal.

Assessing how disruption of the methanogenic community and their syntrophic relationships in tidal freshwater marshes via saltwater intrusion may affect CH₄ emissions

Types of data produced

The field site soil redox and salinity will be measured prior to constructing microcosms. After the microcosms are constructed, methane flux, carbon dioxide flux, butyrate degradation rates, sulfate concentrations, organic matter content, C:N ratio and redox data will be collected from each microcosm during the experiment. Gas flux measurements will be determined using gas chromatography, sulfate and butyrate concentrations will be determined using ion chromatography, organic matter content will be determined gravimetrically, C:N ratios will be determined using a Perkin Elmer Series II CHNS/O Analyzer 2400, and redox and salinity will be measured using a salinity redox probe. In addition total DNA/RNA will be extracted from each microcosm and the 16s gene sequence of both bacteria and methanogens will determined using Illumina sequencing. Data other than the 16s gene sequence data, all other data will be entered into excel and stored using XML formats and/or CSV formats. The environmental data will be analyzed by R and python using ANOVA, regressions analysis, t-tests, and correlated to sequence data using multivariate statistics (e.g. mantel tests). Using multiple reps and including blanks for every variable measured will maintain quality control.

Data and metadata standards

Environmental data, gas flux data, and butyrate degradation data will be entered into excel and stored as both EML and CSV for ease of use in the statistical program R and python. Sequence data from Illumina sequencing will be stored in FASTQ format. The data folder will contain a readme folder the indicates the methods used to collect data on each variable, and the units for the data as well as any labeling schemes. Sequencing conditions (e.g. primer concentration and illumina protocol) will be included in the readme file)

Policies for access and sharing

The environmental data, gas flux measurements, butyrate degradation rates and genetic sequence data will be made available after the resulting manuscript and publication. Until then data will be stored on the VCU microbial ecology R drive and on the Franklin lab Dropbox. All raw data will be made available to Rice Center and if deemed appropriate will be stored on the University's Rice Rivers Center Google Drive. There should be no fee to access the data. Data may also be hosted on a website such as <http://datadryad.org/>, however the data will always be accessible by e-mailing either Dr. Rima Franklin (rbfranklin@vcu.edu) or myself, David Berrier (berrierdj@vcu.edu).

Policies for re-use, distribution

There should be no restrictions on access or reuse of the data. The data will be made available after the publication or two years, whichever ever is first.

Data storage & preservation

Data will be on the Franklin Lab Dropbox and the VCU microbial ecology R drive for long-term storage. The storage of the data on these two different drives in addition to storage on the Rice Center google drive will insure that the data will be preserved. The data will be preserved indefinitely as part of the Franklin lab's database.